

SILICONE FOAM

DRESSING

A HIGHLY ABSORBENT FOAM DRESSING WITH A SOFT SILICONE CONTACT LAYER AND BORDER

Medstock Silicone Foam with Border is a highly absorbent, 5-layer foam dressing that is gentle and conformable to fragile wound tissue. Its perforated silicone skin contact layer provides gentle adhesion to surrounding tissue without sticking to the wound bed.

Key Functions and Features

- + Highly absorbent
- Silicone contact layer seals the wound edges reducing risk of maceration
- + Soft, elastic and comfortable on the skin
- Provides cushioning for wound protection
- + Protective barrier against water and bacteria
- + Gentle and secure adherence
- Atraumatic on dressing removal
 Maintains a moist wound environment to promote healing

Indications for Use


- + Suitable for wounds with moderate to high exudate
- + Skin tear management
- + Venous and arterial leg ulcers
- + Diabetic ulcers
- + Partial thickness burns
- + Surgical wounds
- + Traumatic wounds
- + Pressure injuries
- + Secondary dressing over cavity wounds

General Advice on Safe Use

- + Ensure wound area is clean and dry before application

 Dressing pad should overlap the wound by at least 2cm onto the dry surrounding skin
- → Can be left in place for up to 7 days but should be changed immediately if 3/4 strikethrough (visible exudate) is apparent
- ★ Can be used as a secondary dressing over hydrocolloid paste, alginate and other primary dressings





Skin Contact Layer: Consists of a silicone gel adhesive which works to minimise the risks of trauma upon dressing removal

Wicking Foam Layer: Polyurethane, wicking foam which initiates vertical absorption of wound exudate away from the wound bed and into the superabsorbent layer

Transfer Layer: A non-woven fabric that ensures exudate is efficiently transferred from the wicking foam to the super-absorbent layer

Superabsorbent Layer: The subsurface layer consists of super-absorbent fibres that absorb and retain exudates away from the wound bed

Surface Layer: Highly breathable, complexion-coloured polyurethane film that acts as a waterproof barrier to bacteria



Contraindications / Precautions

- + Arterial bleeds and heavily bleeding wounds
- Do not use Silicone Foam with Border with oxidising agents such as hypochlorite solutions (e.g. EUSOL) or hydrogen peroxide
- + Do not cut Medstock Silicone Foam with Border
- + If reddening or sensitisation occurs, discontinue use and consult a physician

Period of Validity

Caro and Hea Symbols

The product shall be valid for 5 years from the sterilisation date, provided that it is kept under the stipulated storage conditions

Storage Conditions

◆ Store in a well-ventilated clean room at or below 35°C with relative humidity less than 80% and no corrosive gas

Packaging

 Each dressing is packaged in a paper/poly closed pouch, then packed into our unique colour coded Medstock boxes



Medstock Silicone Foam with Border is considered the ideal dressing for skin tears and for use on elderly patients

Care and Ose Symbols						
Read Instructions for us	e. 🕲 Sin	igle use only.	NATEX	Product is latex free.	STERILE EO	Sterilised using ethylene oxide.
Keep dry.	* Sto	ore out of direct sunlight.	®	Do not use if packaging is broken or damaged.		
Product Code			Size			Pcs/Box
PHA7575SF		7.5cm >	7.5cm x 7.5cm		2pcs/pack	
PHA1010SF		10cm x 10cm				2pcs/pack